European Journal of Pharmacology, 242 (1993) 319–321
© 1993 Elsevier Science Publishers B.V. All rights reserved 0014-2999/93/$06.00

Author index

Abiru, T., Watanabe, Y., Kamata, K. and Kasuya, Y.
Simultaneous measurement of vasodilation and changes in cyclic nucleotides in the perfused mesenteric arterial bed of the rat (242) 15
Almotrefi, A.A., see Dzimiri, N. (242) 113
Annies, R., see Löscher, W. (242) 263

Barclay, P.L., see Scott, J.M. (242) 91
Barnes, N.M., see Steward, L.J. (242) 137
Barthelmebs, M., see Michel, B. (242) 237
Beer, M.S. and Middlemiss, D.N.
Serotonin-5-O-carboxymethyl-glycyl[^{125}I]tyrosinamide labels the 5-HT$_{1D\beta}$ receptor subtype in human cortex (242) 195
Biegon, A., see Jagust, W.J. (242) 189
Bisaga, A., Krzascik, P., Jankowska, E., Palejko, W., Kostowski, W. and Danysz, W.
Effect of glutamate receptor antagonists on N-methyl-D-aspartate- and (S)-α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid-induced convulsant effects in mice and rats (242) 213
Bolaños-Jiménez, F., Manhães de Castro, R. and Fillion, G.
Antagonism by citalopram and tianeptine of presynaptic 5-HT$_{1B}$ heteroreceptors inhibiting acetylcholine release (242) 1
Bolwig, T.G., see Glenthøj, B. (242) 283
Bouysset, F., see Vayssettes-Courchay, C. (242) 37
Brennan, K.M., see Jagust, W.J. (242) 189
Bufton, K.E., see Steward, L.J. (242) 137

Campbell, P.M., see Forster, C. (242) 119
Carlsson, A., see Waters, N. (242) 151
Comtois, A., Light, P., Renaud, J.M. and Kong, M.
Tolbutamide, but not glyburide, affects the excitability and contractility of unfatigued frog sartorius muscle (242) 65
Curzon, G., see Gibson, E.L. (242) 83
Członkowski, A., Stein, C. and Herz, A.
Peripheral mechanisms of opioid antinociception in inflammation: involvement of cytokines (242) 229

Danysz, W., see Bisaga, A. (242) 213
Davies, W.E., see Steward, L.J. (242) 137
Davis, B., see Jones, C.A. (242) 183
Dehring, D.J., see Nelson, S.H. (242) 255
Doods, H., Entzeroth, M., Ziegler, H., Schiavi, G., Engel, W., Mihm, G., Rudolf, K. and Eberlein, W.
Characterization of BIBN 99: a lipophilic and selective muscarinic M$_2$ receptor antagonist (242) 23
Dzimiri, N. and Almotrefi, A.A.
Actions of amiodarone on mitochondrial ATPase and lactate dehydrogenase activities in guinea pig heart preparations (242) 113

Eberlein, W., see Doods, H. (242) 23
Eberling, J.L., see Jagust, W.J. (242) 189
Ellis, E.F., see Uphouse, L.A. (242) 53
Embrey, J.P., see Uphouse, L.A. (242) 53
Enas, J.D., see Jagust, W.J. (242) 189

Engel, W., see Doods, H. (242) 23
Entzeroth, M., see Doods, H. (242) 23

Ferrer, J.M.R., see Sabater, R. (242) 205
Fillion, G., see Bolaños-Jiménez, F. (242) 1
Fleetwood-Walker, S.M., Parker, R.M.C., Munro, F.E., Young, M.R., Hope, P.J. and Mitchell, R.
Evidence for a role of tachykinin NK$_2$ receptors in mediating brief nociceptive inputs to rat dorsal horn (laminae III-V) neurons (242) 173
Fölsch, U.R., see Höcker, M. (242) 105
Forster, C. and Campbell, P.M.
Nifedipine inhibits responses to α-adrenoceptor stimulation in canine blood vessels: impact of heart failure (242) 119
Fujita, K., Iijima, N., Suzuki, S. and Nakazawa, K.
Effects of clonidine on desipramine- and desmethylclomipramine-induced 3-methoxy-4-hydroxyphenylglycol levels in rat brain (242) 301

Gessa, G.L., see Rossetti, Z.L. (242) 313
Gibson, E.L., Kennedy, A.J. and Curzon, G.
d-Fenfluramine- and d-norfenfluramine-induced hypophagia: differential mechanisms and involvement of postsynaptic 5-HT receptors (242) 83
Gilmore, J.H., see Watts, V.J. (242) 165
Glenthøj, B., Bolwig, T.G. and Hemmingsen, R.
Effects of chronic discontinuous and continuous treatment of rats with a dopamine D$_1$ receptor antagonist (NNC-756) (242) 283
Grégoire, L., see Hamel, E. (242) 75
Grima, M., see Michel, B. (242) 237
Gutkowska, J., Strick, D.M., Pan, L. and McCann, S.M.
Effect of morphine on urine output: possible role of atrial natriuretic factor (242) 7

Hamel, E., Grégoire, L. and Lau, B.
5-HT$_1$ receptors mediating contraction in bovine cerebral arteries: a model for human cerebrovascular '5-HT$_{1D\beta}$' receptors (242) 75
Hanrahan, S.M., see Jagust, W.J. (242) 189
Hassinen, I.E., see Uusimaa, P.A. (242) 199
Hemmingsen, R., see Glenthøj, B. (242) 283
Herz, A., see Członkowski, A. (242) 229
Higuchi, M., Uezu, K. and Sakanashi, M.
Ex vivo effect of insulin on normal and diabetic rat hearts hypoperfused with norepinephrine (242) 293
Hmaidan, Y., see Rossetti, Z.L. (242) 313
Höcker, M., Hughes, J., Fölsch, U.R. and Schmidt, W.E.
PD 135158, a CCK$_B$/gastrin receptor antagonist, stimulates rat pancreatic enzyme secretion as a CCK$_A$ receptor agonist (242) 105
Hodgson, W.C., see Sikorski, B.W. (242) 275
Hönack, D., see Löscher, W. (242) 263
Hönack, D., see Löscher, W. (242) 99
Hope, P.J., see Fleetwood-Walker, S.M. (242) 173
Hopkins, P.C., see Steward, L.J. (242) 137
Hughes, J., see Höcker, M. (242) 105

Iijima, N., see Fujita, K. (242) 301
Imbs, J.-L., see Michel, B. (242) 237
Ito, M., see Suga, T. (242) 129
Itoh, H., see Suga, T. (242) 129
Iwamoto, Y., see Kamei, J. (242) 209

Jagust, W.J., Eberling, J.L., Roberts, J.A., Brennan, K.M., Hanrahan, S.M., VanBrocklin, H., Enas, J.D., Biegon, A. and Mathis, C.A.
In vivo imaging of the 5-hydroxytryptamine reuptake site in primate brain using single photon emission computed tomography and [^{123}I]5-iodo-6-nitroquipazine (242) 189
Jankowska, E., see Bisaga, A. (242) 213
Johnson, R.L., see Nelson, S.H. (242) 255
Jones, C.A., Zempléni, É., Davis, B. and Reynolds, G.P.
Glutamate stimulates dopamine release from cortical and limbic rat brain in vitro (242) 183
Junien, J.-L., see Roman, F.J. (242) 305

Kamata, K., see Abiru, T. (242) 15
Kamei, J., Iwamoto, Y., Misawa, M. and Kasuya, Y.
Effects of rimcazole, a specific antagonist of σ sites, on the antitussive effects of non-narcotic antitussive drugs (242) 209
Kasuya, Y., see Abiru, T. (242) 15
Kasuya, Y., see Kamei, J. (242) 209
Katsumata, K., see Sakurada, T. (242) 47
Kennedy, A.J., see Gibson, E.L. (242) 83
King, R.G., see Sikorski, B.W. (242) 275
Kisara, K., see Sakurada, T. (242) 47
Kiviluoma, K.T., see Uusimaa, P.A. (242) 199
Klitgaard, H., Knutsen, L.J.S. and Thomsen, C.
Contrasting effects of adenosine A_1 and A_2 receptor ligands in different chemoconvulsive rodent models (242) 221
Knutsen, L.J.S., see Klitgaard, H. (242) 221
Kong, M., see Comtois, A. (242) 65
Konishi, T., see Suga, T. (242) 129
Kostowski, W., see Bisaga, A. (242) 213
Krzascik, P., see Bisaga, A. (242) 213
Kusagawa, M., see Suga, T. (242) 129

Lagerkvist, S., see Waters, N. (242) 151
Lai, M., see Rossetti, Z.L. (242) 313
Lamb, F.S., see Watts, S.W. (242) 245
Land, T., see Ögren, S.O. (242) 59
Langel, Ü., see Ögren, S.O. (242) 59
Lau, B., see Hamel, E. (242) 75
Laubie, M., see Vayssettes-Courchay, C. (242) 37
Lawler, C.P., see Watts, V.J. (242) 165
Light, P., see Comtois, A. (242) 65
Löfberg, L., see Waters, N. (242) 151
Löscher, W. and Hönack, D.
Effects of the novel 5-HT_{1A} receptor antagonist, (+)-WAY 100135, on stereotyped behaviour induced by the NMDA receptor antagonist dizocilpine in rats (242) 99
Löscher, W., Annies, R. and Hönack, D.
Comparison of competitive and uncompetitive NMDA receptor antagonists with regard to monoaminergic neuronal activity and behavioural effects in rats (242) 263
Löscher, W., see Richter, A. (242) 309

Mailman, R.B., see Watts, V.J. (242) 165
Manhães de Castro, R., see Bolaños-Jiménez, F. (242) 1
Manome, Y., see Sakurada, T. (242) 47
Martin, B., see Roman, F.J. (242) 305
Mathis, C.A., see Jagust, W.J. (242) 189
Mayhan, W.G.
In vivo reactivity of resistance arterioles to activation of ATP-sensitive K^+ channels (242) 109
McCann, S.M., see Gutkowska, J. (242) 7
Michel, B., Stephan, D., Grima, M., Barthelmebs, M. and Imbs, J.-L.
Effects of one-hour and one-week treatment with ramipril on plasma and renal brush border angiotensin converting enzyme in the rat (242) 237
Middlemiss, D.N., see Beer, M.S. (242) 195
Mihm, G., see Doods, H. (242) 23
Misawa, M., see Kamei, J. (242) 209
Mitchell, R., see Fleetwood-Walker, S.M. (242) 173
Munro, F.E., see Fleetwood-Walker, S.M. (242) 173
Myers, J.H., see Watts, S.W. (242) 245

Nakano, T., see Suga, T. (242) 129
Nakazawa, K., see Fujita, K. (242) 301
Nelson, S.H., Suresh, M.S., Dehring, D.J. and Johnson, R.L.
Relaxation by calcitonin gene-related peptide may involve activation of K^+ channels in the human uterine artery (242) 255
Nichols, D.E., see Watts, V.J. (242) 165

Ögren, S.O., Pramanik, A., Land, T. and Langel, Ü.
Differential effects of the putative galanin receptor antagonists M15 and M35 on striatal acetylcholine release (242) 59
Ohba, M., see Sakurada, T. (242) 47

Palejko, W., see Bisaga, A. (242) 213
Pan, L., see Gutkowska, J. (242) 7
Parker, R.M.C., see Fleetwood-Walker, S.M. (242) 173
Peuhkurinen, K.J., see Uusimaa, P.A. (242) 199
Piercey, M., see Waters, N. (242) 151
Portoghese, P.S., see Takemori, A.E. (242) 145
Pramanik, A., see Ögren, S.O. (242) 59

Renaud, J.M., see Comtois, A. (242) 65
Reynolds, G.P., see Jones, C.A. (242) 183
Richter, A. and Löscher, W.
The atypical neuroleptic, clozapine, exerts antidystonic activity in a mutant hamster model. Comparison with haloperidol (242) 309
Roberts, J.A., see Jagust, W.J. (242) 189
Roman, F.J., Martin, B. and Junien, J.-L.
In vivo interaction of neuropeptide Y and peptide YY with σ receptor sites in the mouse brain (242) 305
Rossetti, Z.L., Lai, M., Hmaidan, Y. and Gessa, G.L.
Depletion of mesolimbic dopamine during behavioral despair: partial reversal by chronic imipramine (242) 313
Rudolf, K., see Doods, H. (242) 23
Ruskoaho, H., see Uusimaa, P.A. (242) 199

Sabater, R., Saez, J.A. and Ferrer, J.M.R.
SCH 23390 decreases self-stimulation of the medial prefrontal cortex in the rat (242) 205
Saez, J.A., see Sabater, R. (242) 205
Sakanashi, M., see Higuchi, M. (242) 293
Sakurada, S., see Sakurada, T. (242) 47
Sakurada, T., Katsumata, K., Manome, Y., Tan-No, K., Sakurada, S., Kisara, K. and Ohba, M.
Antinociceptive effects in the formalin and capsaicin tests after intrathecal administration of substance P analogues in mice (242) 47
Schiavi, G., see Doods, H. (242) 23
Schmidt, W.E., see Höcker, M. (242) 105
Schoemaker, H.
[^{3}H]7-OH-DPAT labels both dopamine D_3 receptors and σ sites in the bovine caudate nucleus (242) R1

Scott, J.M., Barclay, P.L. and Shepperson, N.B.
Renal effects of neutral endopeptidase inhibition in euvolemic and hypervolemic rats (242) 91
Shepperson, N.B., see Scott, J.M. (242) 91
Shimomura, A., see Suga, T. (242) 129
Sikorski, B.W., Hodgson, W.C. and King, R.G.
Effects of haemoglobin and N-nitro-L-arginine on constrictor and dilator responses of aortic rings from streptozotocin diabetic rats (242) 275
Silverman, P.B.
On-off effects of dopamine receptor agonists in the hemi-parkinsonian rat (242) 31
Southerland, S.B., see Watts, V.J. (242) 165
Stein, C., see Członkowski, A. (242) 229
Stephan, D., see Michel, B. (242) 237
Steward, L.J., Bufton, K.E., Hopkins, P.C., Davies, W.E. and Barnes, N.M.
Reduced levels of 5-HT_3 receptor recognition sites in the putamen of patients with Huntington's disease (242) 137
Strick, D.M., see Gutkowska, J. (242) 7
Suga, T., Itoh, H., Shimomura, A., Kusagawa, M., Ito, M., Takase, K., Konishi, T. and Nakano, T.
Comparison of the effects of various vasodilators on the rat portal vein and mesenteric artery (242) 129
Suresh, M.S., see Nelson, S.H. (242) 255
Suzuki, S., see Fujita, K. (242) 301

Takase, K., see Suga, T. (242) 129
Takemori, A.E. and Portoghese, P.S.
Enkephalin antinociception in mice is mediated by δ_1-and δ_2-opioid receptors in the brain and spinal cord, respectively (242) 145
Tan-No, K., see Sakurada, T. (242) 47
Thomsen, C., see Klitgaard, H. (242) 221
Tóth, M., see Uusimaa, P.A. (242) 199
Traub, O., see Watts, S.W. (242) 245

Uezu, K., see Higuchi, M. (242) 293
Uphouse, L.A., Welch, S.P., Ward, C.R., Ellis, E.F. and Embrey, J.P.
Antinociceptive activity of intrathecal ketorolac is blocked by the κ-opioid receptor antagonist, nor-binaltorphimine (242) 53
Uusimaa, P.A., Kiviluoma, K.T., Tóth, M., Hassinen, I.E., Peuhkurinen, K.J. and Ruskoaho, H.
Adenosine inhibits the release of atrial natriuretic peptide from the perfused rat heart (242) 199

VanBrocklin, H., see Jagust, W.J. (242) 189
Vayssettes-Courchay, C., Bouysset, F., Verbeuren, T.J. and Laubie, M.
Role of the nucleus tractus solitarii and the rostral depressive area in the sympatholytic effect of 8-hydroxy-2-(di-n-propylamino)tetralin in the cat (242) 37
Verbeuren, T.J., see Vayssettes-Courchay, C. (242) 37

Ward, C.R., see Uphouse, L.A. (242) 53
Watanabe, Y., see Abiru, T. (242) 15
Waters, N., Lagerkvist, S., Löfberg, L., Piercey, M. and Carlsson, A.
The dopamine D_3 receptor and autoreceptor preferring antagonists (+)-AJ76 and (+)-UH232; a microdialysis study (242) 151
Watts, S.W., Traub, O., Lamb, F.S., Myers, J.H. and Webb, R.C.
Effect of ramipril on α-adrenoceptor-mediated oscillatory contractions in tail artery of hypertensive rats (242) 245
Watts, V.J., Lawler, C.P., Gilmore, J.H., Southerland, S.B., Nichols, D.E. and Mailman, R.B.
Dopamine D_1 receptors: efficacy of full (dihydrexidine) vs. partial (SKF38393) agonists in primates vs. rodents (242) 165
Webb, R.C., see Watts, S.W. (242) 245
Welch, S.P., see Uphouse, L.A. (242) 53

Young, M.R., see Fleetwood-Walker, S.M. (242) 173

Zempléni, É., see Jones, C.A. (242) 183
Ziegler, H., see Doods, H. (242) 23